

To make your breathing better, you MUST take your medicine as explained below. Following these instructions puts more of the medicine into your lungs. This will open your air passages and help you breathe easier and feel better. You need to ask your health-care provider or pharmacist how many puffs of medicine your metered-dose inhaler (MDI) has when it is full. You need to keep track of how many puffs of medicine you take everyday, so you can have your MDI refilled before you run out of medicine. Before using your MDI, please read the priming or preparing instructions. Your MDI should be cleaned once a week. See the instructions on cleaning your MDI.



Take cap off MDI. Check for and remove any dust, lint, or other objects. Shake MDI well.



Sit up straight or stand up.




Hold your MDI two finger widths away from your lips.



Breathe out all the way.

The American College of Chest Physicians is the leading resource for the improvement of cardiopulmonary health and critical care worldwide. Its mission is to promote the prevention and treatment of diseases of the chest through leadership, education, research, and communication.

This publication's content contains general information, is not intended to be and is not complete, is not medical advice, and does not replace professional medical care and physician advice, which always should be sought for any specific condition. The American College of Chest Physicians and its officers, regents, executive committee, members, and employees specifically disclaim all responsibility for any liability, damages (actual or consequential), loss, or risk, personal or otherwise, based on any legal theory whatsoever, alleged to have been incurred as a result, directly or indirectly, of the use of any of the material herein.





American College of Chest Physicians 3300 Dundee Road, Northbrook, IL 60062 (847) 498-1400 phone (847) 498-5460 fax www.chestnet.org





As you begin to **BREATHE IN SLOWLY, PRESS DOWN ON THE MDI**, as shown in this picture. Keep breathing in until your lungs are completely full.



HOLD your breath for 10 seconds. If you cannot hold your breath for 10 seconds, hold your breath as long as you can.

7	
	11 12 1
	8 4
	765


If you need to take another puff of medicine, wait 1 minute. After 1 minute, repeat steps 2-6.



Rinse your mouth out after you take your last puff of medicine. Make sure you spit the water out; do not swallow it. *Rinsing is only necessary if the medicine you just took was a corticosteroid, such as Flovent®, Beclovent®, Vanceril®, Aerobid®, or Azmacort®.Recap the MDI.*

The American College of Chest Physicians is the leading resource for the improvement of cardiopulmonary health and critical care worldwide. Its mission is to promote the prevention and treatment of diseases of the chest through leadership, education, research, and communication.

This publication's content contains general information, is not intended to be and is not complete, is not medical advice, and does not replace professional medical care and physician advice, which always should be sought for any specific condition. The American College of Chest Physicians and its officers, regents, executive committee, members, and employees specifically disclaim all responsibility for any liability, damages (actual or consequential), loss, or risk, personal or otherwise, based on any legal theory whatsoever, alleged to have been incurred as a result, directly or indirectly, of the use of any of the material herein.




Priming Your Metered-Dose Inhaler (MDI)

Preparing your MDI will need to be done for every new MDI before it is used or if it has been a long time since you last used your MDI. There are two types of MDIs. You need to ask your health-care provider or pharmacist if your MDI is a HFA or a CFC. The type of MDI you have will dictate how you prepare your MDI before use.

HFA MDIs: If your MDI is new or has not been used in 2 weeks, shake the MDI very well, then press down on the MDI four times, wasting four puffs into the air. After this is done, you are ready to take your medicine. Follow the instructions below to learn how to take your medicine the correct way. See instructions on using your MDI (separate handout).







CFC MDIs: If your MDI is new or has not been used in 12 hours, shake the MDI very well, then press down on the MDI once, wasting one puff of medicine into the air. After this is done, you are ready to take your medicine. Follow the instructions below to learn how to take your medicine the correct way. See instructions on using your MDI (separate handout).

Cleaning Your MDI



Take metal canister out of plastic container.



Rinse plastic container with warm water at least once a week.



Let plastic container air dry.



Replace metal canister in plastic container.

The American College of Chest Physicians is the leading resource for the improvement of cardiopulmonary health and critical care worldwide. Its mission is to promote the prevention and treatment of diseases of the chest through leadership, education, research, and communication.

This publication's content contains general information, is not intended to be and is not complete, is not medical advice, and does not replace professional medical care and physician advice, which always should be sought for any specific condition. The American College of Chest Physicians and its officers, regents, executive committee, members, and employees specifically disclaim all responsibility for any liability, damages (actual or consequential), loss, or risk, personal or otherwise, based on any legal theory whatsoever, alleged to have been incurred as a result, directly or indirectly, of the use of any of the material herein.